



Registration Ref.	#121 (30/5/2008)
Cultivar Name	<b><i>P.</i> 'Aninha'</b>
Originator	Margarete M. Souza (Brazil)
Parentage	<i>P. palmeri</i> var. <i>sublanceolata</i> (♀) × <i>P. foetida</i> var. <i>foetida</i> (♂)
Description	Vine pilose, stem cylindrical; internodes 6.2 – 11.0cm; flowers pale pink, 6 - 7cm in diameter, reflexing at anthesis (before midday); petals 2.5 - 2.9cm long, 0.9 - 1.2cm wide, pale pink on both surfaces, apex ovate-lanceolate; sepals, 2.6 - 3.1cm long, 0.9 - 1.2cm wide, adaxial surface pale pink, abaxial surface green, apex ovate-lanceolate; corona filaments in 4 series, outermost series 1.0 - 1.4cm long, second series 1.1 - 1.5cm long, filiform, reddish/purple base, white, purple/white banding to apex; leaves three-lobed, membranaceous, hairy on both sides, 8.7 - 11.6cm long, 6.6 - 8.3cm wide; peduncles stout, 4.3 - 6.7cm; stamen filaments pale green; anthers adaxial surface pale green with yellow edges; style slender, pale green; stigma olive green; fruits pendulous, ripening to scarlet. Flowering all year in Brazil. Although based on a different variety of <i>P. foetida</i> , this cultivar is the reciprocal cross of <i>P.</i> 'Pink Jewel' at the species level.