



Registration Ref.	#148 (11/8/2009)
Cultivar Name	<i>P. 'Murac'</i>
Originator	Franz Düllli (Switzerland) and Axel Frank (Germany)
Parentage	<i>P. mucronata</i> (♀) × <i>P. racemosa</i> (♂)
Description	Stem round; vine vigourous, floriferous; flowers up to 11cm in diameter, unscented, reflexing, anthesis in afternoon, with flowers closing next morning; corona filaments 4cm in diameter, two outer rows greenish/white banded at the centre, white to apex; two shorter inner rows greenish-white; sepals 4.6 x 1.1cm, adaxial surface pink/white, abaxial surface burgundy-red, slightly keeled; petals 4.5 x 0.9cm, abaxial surface similar to, but slightly paler than sepals, abaxial surface white; peduncle 5-8cm; bracts red, oval, 2.5-2.8 x 1.6-2.0cm with awn, 2cm below bud; leaves unlobed, oblong/oval, pale green, red bordered, abaxial surface red-veined, 8.1-9.5 x 5.2-5.7cm; stipules 2.4-2.6 x 1.3-1.5cm with 0.1cm awn, inside light green with red-borders, outside red-veined, on young shoots the stipules are red on the outer surface; petioles 2.0-3.2cm, with a pair of glands. This hybrid was created to investigate the parentage of <i>P. 'Kew Gardens'</i> [see <i>Passiflorunde</i> 15(1), 2-4, 2007]. The plant was tested for two years and overwintered at 10C. The name is an acronym derived from the names of the parents.